

NAC (N-Acetyl-L-Cysteine)

NAC — N-Acetyl-L-Cysteine, an acetylated form of cysteine, is readily absorbed to immediately provide an efficient precursor of cysteine for continued glutathione (GSH) synthesis.


The amino acid cysteine in its pre-acetylated form is known as N-Acetyl-L-Cysteine. It is essential for glutathione synthesis as it serves as the precursor to glutathione intracellular synthesis.^(1,2) Glutathione is a critical component of the immune system, as it is an important intracellular water-soluble antioxidant. As part of this function, glutathione plays a vital role as the cofactor for the antioxidant enzymes glutathione peroxidase and glutathione transferase. Glutathione is also important to the liver, as the liver utilizes it for drug detoxification. As a metabolite of cysteine, **NAC** plays an important role in detoxification. As a more stable form of L-Cysteine, **NAC** functions in protecting cells against oxidative stress.⁽³⁾ Additionally, **NAC** contributes to important bodily reactions, making it a part of a defensive mechanism toward potential carcinogens as well as against DNA damage. The role of **NAC** in acetaminophen toxicity is widely documented, however, **NAC** contributes to other protective mechanisms, including drug conjugation and excretion, peroxides and free radical destruction, maintenance of the redox state of NADPH-NADP, acid-base balance, and the synthesis of sulfated compounds.⁽⁴⁾ **NAC** also plays a role in the development of the central nervous system.⁽⁵⁾

The protective method of **NAC** has been attributed to its:

- Nucleophilicity, antioxidant activity
- Modulation of the metabolism
- Effects in the mitochondria

- Decrease in the biologically effective dose of carcinogens
- Modulation of DNA repair
- Inhibition of genotoxicity and cell transformation
- Modulation of gene expression and signal transduction pathways
- Regulation of cell survival and apoptosis
- Anti-inflammatory activity and anti-angiogenic activity
- Immunological effects
- Inhibition of cellular dysfunction
- Influence on cell cycle progression
- Inhibition of hypercellular growth
- Inhibition of invasion and abnormal cell proliferation
- Ability to block NF-kappaB activation⁽⁶⁾
- Protection against the adverse effects of chemopreventive and chemotherapeutic agents⁽¹⁾

There have been numerous studies using **NAC**, with COPD, particularly amongst patients who smoke. Utilizing **NAC** in smokers with chronic bronchitis showed a significant reduction in the number of positive bacterial cultures in these patients.⁽⁷⁾ An additional six-month study demonstrating the effects of **NAC** on chronic bronchitis indicated a significant reduction in sick-leave days for the **NAC** group, compared to placebo. Another study utilizing **NAC** indicated potential benefits for chronic bronchitis, resulting in a significantly lower exacerbation rate in patients administered **NAC**,



(800) 231-5777

6801 Biotics Research Drive • Rosenberg, TX 77471
biotics@bioticsresearch.com • www.bioticsresearch.com

These statements have not been evaluated by the Food and Drug Administration. These products are not intended to diagnose, treat, cure, or prevent any disease.

compared to placebo.⁽⁸⁾ In the group administered **NAC**, 40% remained free from exacerbations, compared to 19% in the placebo group. The study also indicated that the sick-leave rate due to exacerbation in the **NAC** group was significantly infrequent.⁽⁹⁾

NAC has also been shown to be radioprotectant against oxidative damage.⁽¹⁰⁾ Additionally, researchers have cited the beneficial use of **NAC** for the improvement of insulin sensitivity in women with polycystic ovary syndrome (PCOS), noting a significant improvement in circulating insulin levels, as well as insulin sensitivity in these patients.⁽¹¹⁾

Since it easily passes through cellular membranes, **NAC** has been shown to be readily absorbed. It is also resistant to enzymatic breakdown and is important for cellular glutathione production⁽¹²⁾

NAC supplies 500mg of N-Acetyl-L-Cysteine per tablet.

References

1. Issels RD, Nagele A, Eckert KG, Wilmanns W. Promotion of cystine uptake and its utilization for glutathione biosynthesis induced by cysteamine and N-acetylcysteine. *Biochem Pharmacol.* 1988 Mar 1;37(5):881-8.
2. Phelps DT, Deneke SM, Daley DL, Fanburg BL. Elevation of glutathione levels in bovine pulmonary artery endothelial cells by N-acetylcysteine. *Am J Respir Cell Mol Biol.* 1992 Sep; 7(3):293-9.
3. **Biochemistry**, Cantarow & Schepartz, 1954
4. De Flora S., Izzotti A, D'Agostini F, Balansky RM. Mechanisms of N-acetylcysteine in the prevention of DNA damage and cancer, with special reference to smoking-related end-points *Carcinogenesis.* 2001 Jul;22(7):999-1013.
5. Martha H. Stipanuk. www.nutrition.Cornell.edu.
6. Pajonk F, Riess K, Sommer A, McBride WH. N-acetyl-L-cysteine inhibits 26S proteasome function: implications for effects on NF-kappaB activation. *Free Radic Biol Med.* 2002 Mar 15;32(6):536-43.
7. Riise GC, Larsson S, Larsson P, Jeansson S, Andersson BA. The intrabronchial microbial flora in chronic bronchitis patients: a target for N-acetylcysteine therapy? *Eur Respir J.* 1994 Jan; 7(1):94-101.

8. Rasmussen, JB & Glennow, C. Reduction in days of illness after long-term treatment with N-acetylcysteine controlled-release tablets in patients with chronic bronchitis. *Eur Respir J.* 1988 Apr;1(4):351-5.
9. Bowman G, Backer U, Larsson S, Melander B, Wahlander L. Oral acetylcysteine reduces exacerbation rate in chronic bronchitis: report of a trial organized by the Swedish Society for Pulmonary Disease. *Eur J Respir Dis.* 1983 Aug; 64(6):405-415.
10. Neal R, Matthews RH, Lutz P, Ercal N. Antioxidant Role of N-Acetyl cysteine isomers following high dose irradiation. *Free Radic Biol Med.* 2003 Mar 15;34(6):689-95.
11. Fulghesu AM, Ciampelli M, Muzj G, Belosi C, Selvaggi L, Ayala GF, Lanzone A. N-acetyl-cysteine treatment improves insulin sensitivity in women with polycystic ovary syndrome. *Fertil Steril.* 2002 Jun;77(6):1128-35.
12. DeCaro L, Ghizzi A, Costa R, Longo A, Ventresca GP, Lodola E. Pharmacokinetics and bioavailability of oral acetylcysteine in healthy volunteers. *Arzneimittelforschung.* 1989 Mar;39(3):382-6.

Supplement Facts

Serving Size: 1 Capsule

	Amount Per Serving
N-Acetyl-L-Cysteine	500 mg*

* Daily Value not established

Other ingredients: Capsule shell (gelatin and water).

This product is gluten and dairy free.

RECOMMENDATION: One (1) capsule each day as a dietary supplement or as otherwise directed by a healthcare professional.

KEEP OUT OF REACH OF CHILDREN

Store in a cool, dry area.
Sealed with an imprinted safety seal for your protection.

Product # 5205 Rev. 12/14

To place your order for **NAC** or for additional information please contact us below.



BIOTICS
RESEARCH
CORPORATION
Utilizing "The Best of Science and Nature"
to Create Superior Nutritional Supplements



(800) 231-5777

6801 Biotics Research Drive • Rosenberg, TX 77471
biotics@bioticsresearch.com • www.bioticsresearch.com

These statements have not been evaluated by the Food and Drug Administration. These products are not intended to diagnose, treat, cure, or prevent any disease.